

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

[www.elsevier.com/locate/jes](http://www.elsevier.com/locate/jes)

**JES**  
JOURNAL OF  
ENVIRONMENTAL  
SCIENCES  
[www.jesc.ac.cn](http://www.jesc.ac.cn)

# Large variability in ambient ozone sensitivity across 19 ethylenediurea-treated Chinese cultivars of soybean is driven by total ascorbate

Lijun Jiang<sup>1,2,3</sup>, Zhaozhong Feng<sup>1,2,3,\*</sup>, Lulu Dai<sup>1,3</sup>, Bo Shang<sup>1,3</sup>, Elena Paoletti<sup>1,4</sup>

1. State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

2. State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese Academy of Sciences, Nanjing 210008, China

3. College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

4. National Research Council, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy

## ARTICLE INFO

### Article history:

Received 8 March 2017

Revised 30 June 2017

Accepted 4 July 2017

Available online 12 July 2017

### Keywords:

Ground-level ozone

Glycine max

Cultivar

Ozone sensitivity

Antioxidants

Gas exchange

Biomass

Total ascorbate

## ABSTRACT

The sensitivity of Chinese soybean cultivars to ambient ozone ( $O_3$ ) in the field is unknown, although soybean is a major staple food in China. Using ethylenediurea (EDU) as an  $O_3$  protectant, we tested the gas exchange, pigments, antioxidants and biomass of 19 cultivars exposed to 28 ppm-hr AOT40 (accumulated  $O_3$  over an hourly concentration threshold of 40 ppb) over the growing season at a field site in China. By comparing the average biomass with and without EDU, we estimated the cultivar-specific sensitivity to  $O_3$  and ranked the cultivars from very tolerant (<10% change) to highly sensitive (>45% change), which helps in choosing the best-suited cultivars for local cultivation. Higher lipid peroxidation and activity of the ascorbate peroxidase enzyme were major responses to  $O_3$  damage, which eventually translated into lower biomass production. The constitutional level of total ascorbate in the leaves was the most important parameter explaining  $O_3$  sensitivity among these cultivars. Surprisingly, the role of stomatal conductance was insignificant. These results will guide future breeding efforts towards more  $O_3$ -tolerant cultivars in China, while strategies for implementing control measures of regional  $O_3$  pollution are being implemented. Overall, these results suggest that present ambient  $O_3$  pollution is a serious concern for soybean in China, which highlights the urgent need for policy-making actions to protect this critical staple food.

© 2017 The Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences.

Published by Elsevier B.V.

## Introduction

Food security is a topical issue nowadays, especially in rapidly expanding China (Yin et al., 2009). China is the fourth largest world producer of soybean (*Glycine max* (L.) Merr.), with 12.2 million tons in 2014 (FAO, 2014). Soybean is a key source

of vegetable protein for humans (Mateosapario et al., 2008). It is one of the most important agricultural crop species and the top legume species worldwide (FAO, 2013).

China is currently suffering from serious surface ozone ( $O_3$ ) pollution, with annual peak averages reaching as high as 60 ppb (Feng et al., 2015) and an increase of about 7% from

\* Corresponding author. E-mail: [fzz@rcees.ac.cn](mailto:fzz@rcees.ac.cn) (Zhaozhong Feng).

2005 to 2010 (Verstraeten et al., 2015). Ozone is one of the most detrimental air pollutants for crops and natural ecosystems (Ainsworth et al., 2012). Soybean ranks among the most O<sub>3</sub>-sensitive agricultural crops (Mills et al., 2007), and such current O<sub>3</sub> concentrations are high enough to cause significant yield losses (Morgan et al., 2003). Projected O<sub>3</sub>-induced soybean yield losses were 9.5%–15% for the year 2030 at the global level (Avnery et al., 2011), and the financial losses for soybean were estimated as 2.0–5.8 billion US dollars annually based on the price in the year 2000 (Osborne et al., 2016). Many experiments in different parts of the world have been carried out to investigate the physiological, growth and yield responses of soybean to O<sub>3</sub> in open-top chambers and under ambient conditions (e.g., Sun et al., 2014; Zhang et al., 2014; Rai et al., 2015). Ozone exposure reduces photosynthesis, stomatal conductance and the leaf chlorophyll content of soybean (Morgan et al., 2003). A SoyFACE study showed a dose-dependent linear decrease in soybean yield and photosynthesis, and altered antioxidant capacity (Betzlberger et al., 2012).

Dose–response studies for a range of crops have revealed that O<sub>3</sub> sensitivity is a heritable trait (Reinert and Eason, 2000) and is highly variable among species and cultivars (Ariyaphanphitak et al., 2005; Mills et al., 2007). Studies on the response of soybean to O<sub>3</sub> in Asia have focused on the growth and yield of individual cultivars (Wahid et al., 2001; Singh et al., 2010; Singh and Agrawal, 2011; Rai et al., 2015). However, Zhang et al. (2014) demonstrated that O<sub>3</sub> sensitivity varied greatly across nine soybean cultivars widely cultivated in Northeast China, through elevated O<sub>3</sub> exposure experiments in open-top chambers. So far, however, there is no available data showing whether current ambient O<sub>3</sub> levels affect the growth and productivity of soybean in China.

The antiozonant ethylenediurea (N-[2-(2-oxo-1-imidazolidinyl)ethyl]-N'-phenylurea, abbreviated as EDU, with formula C<sub>4</sub>H<sub>10</sub>N<sub>4</sub>O<sub>2</sub>), first described by Carnahan et al. (1978), is a well-known antiozonant chemical (Paoletti et al., 2009; Feng et al., 2010; Manning et al., 2011; Agathokleous et al., 2016a), able to prevent O<sub>3</sub> injury, especially visible foliar O<sub>3</sub> injury as well as growth reduction in agricultural and horticultural crops and forest trees, by stimulating the antioxidant defense (Tiwari et al., 2005; Elagöz and Manning, 2005; Szantoi et al., 2007; Paoletti et al., 2007; Feng et al., 2010; Rai et al., 2015). A meta-analysis suggested that the antiozonant activity of EDU is biochemical rather than biophysical (Feng et al., 2010), but conclusive proof of the detailed basis for the protective action has not been confirmed. Recent results showed that EDU does not have side-effects on growth and is not toxic to plants at the concentrations required for O<sub>3</sub> protection (Agathokleous et al., 2016a). As a reliable, low-cost, and low-technology tool, EDU has great potential for assessing the effects of ambient O<sub>3</sub> on vegetation (Singh et al., 2014; Agathokleous et al., 2016b, 2016c).

We used EDU as a tool for assessing: (1) the relative sensitivity to ambient O<sub>3</sub> exposure in 19 soybean cultivars widely cultivated in China by using biomass as the response indicator, (2) whether these cultivars differ in their physiological and biochemical responses to O<sub>3</sub> (gas exchange, pigments, antioxidants), and (3) which parameters are the most important as predictors of O<sub>3</sub>-sensitivity in these cultivars. This knowledge will help in cultivating the most O<sub>3</sub>-tolerant cultivars in the areas at higher risk and breeding for more and more O<sub>3</sub>-tolerant cultivars.

## 1. Materials and methods

### 1.1. Experimental conditions

The experiment was conducted under natural field conditions from June to October, 2015, at a suburban area of Beijing city, Changping District, 40°19' N, 116°13' E and 43.5 m a.s.l. (above sea level). The site is about 52 km from the city center. Mean monthly minimum and maximum temperatures were –3.1 °C (January) and 26.7 °C (July). The mean yearly precipitation was 550 mm and almost 60% of rain occurred in July and August.

Meteorological variables (air temperature and precipitation) were recorded by a portable automatic weather station (HOBO-U30, USA). The concentration of O<sub>3</sub> was continuously monitored using an ultraviolet (UV)-absorption O<sub>3</sub> analyzer (Model 49i, Thermo Scientific, USA). The monitor was calibrated by a 49i-PS calibrator (49i-PS, Thermo Scientific, USA) before the experiment and once a month during the experiment. Exceedances above 40 ppb were accumulated to calculate the exposure index AOT40 (accumulated O<sub>3</sub> over an hourly concentration threshold of 40 ppb) according to Mills et al. (2007). The distribution of hourly O<sub>3</sub> concentrations across 10 ppb classes of exposure and daily 8-hr means was calculated from 9:00 to 17:00 solar time.

The seeds of 19 soybean cultivars (*Glycine max* (L.) Merr.) were obtained from the Institute of Crop Science of Chinese Academy of Agricultural Sciences. The cultivars are widely planted in North China, have similar growing periods (110–130 days), and had not been tested for O<sub>3</sub> sensitivity previously. The agronomic characteristics of these cultivars are listed in Table 1. The soybean seeds were sown on the 10th of June and sprouted out of the earth on the 20th of June, 2015.

After measuring the physiological and biochemical parameters at two months after germination (23rd August), harvest was carried out at the very end of the growing season (8th October). Due to rainy days at the time of flowering (22nd July to 10th August) (Fig. 1), the plants did not produce seeds. No soybean yield occurred in the entire region in 2015. Therefore, the present paper shows only the results of biomass.

In this experiment, there were 7 plots and each plot occupied 65 m<sup>2</sup>. For every plot, there were 19 lines (5 m in length for each line) i.e., one line per cultivar distributed at random. The basic physical and chemical properties of soil were as follows: organic C, 17.4 g/kg; total N, 0.9 mg/kg; available P, 38.1 mg/kg; available K, 102.1 mg/kg and pH of 8.3.

### 1.2. EDU application

Among different concentrations of EDU, 450 ppm of EDU was used in this study as it was found to effectively protect different plant species from O<sub>3</sub> (Paoletti et al., 2009; Feng et al., 2010; Manning et al., 2011). For instance, foliar applications of EDU at 450 ppm significantly alleviated snap bean foliar injury, and increased the photosynthesis rate, seed and pod weights in O<sub>3</sub>-sensitive genotypes (Yuan et al., 2015). EDU powder (100% available ingredient) was dissolved in warm water. Three plots were sprayed with water and four plots were sprayed with EDU. The entire foliage of each plant was sprayed until the drip point before sunrise each time. The EDU treatments started from the

**Table 1 – Characteristics of the 19 soybean cultivars used in the study.**

| Cultivars    | Cultivars (abbreviation) | Year of release | Maturity (days) | Male parent  | Female parent     | Leaf shape |
|--------------|--------------------------|-----------------|-----------------|--------------|-------------------|------------|
| Jidou12      | J12                      | 2002            | 100             | Oil 83-14    | Jinda7826         | Ovoid      |
| ZhongHuang13 | ZH13                     | 2001            | 105             | Yudou8       | Zhongzuo90052-76  | Ovoid      |
| ZhongHuang20 | ZH20                     | 2003            | 100             | Yi-2         | Hobbit            | Ovoid      |
| ZhongHuang39 | ZH39                     | 2006            | 100             | Zhongpin661  | Zhonghuang14      | Ovoid      |
| ZhongHuang40 | ZH40                     | 2007            | 104             | Jindou6      | Yudou12           | Ovoid      |
| ZhongHuang41 | ZH41                     | 2009            | 108             | Kefeng14     | Kexin3            | Ovoid      |
| ZhongHuang42 | ZH42                     | 2007            | 116             | Yochu4       | Jindou33          | Ovoid      |
| ZhongHuang43 | ZH43                     | 2006            | 101             | Jidou7       | Xinke3            | Ovoid      |
| ZhongHuang44 | ZH44                     | 2009            | 107             | Kefeng14     | Kexin3            | Ovoid      |
| ZhongHuang48 | ZH48                     | 2009            | 108             | Kefeng14     | Kexin3            | Ovoid      |
| ZhongHuang49 | ZH49                     | 2009            | 106             | Kefeng14     | Kexin3            | Ovoid      |
| ZhongHuang50 | ZH50                     | 2010            | 106             | Zhonghuang13 | Zhongpin661       | Ovoid      |
| ZhongHuang62 | ZH62                     | 2011            | 100             | Zhonghuang25 | Xindou1           | Lanceolate |
| ZhongHuang66 | ZH66                     | 2014            | 112             | Zhongpin661  | Cheng9039-2-4-3-1 | Ovoid      |
| ZhongHuang69 | ZH69                     | 2012            | 121             | Kefeng14     | Kexin3            | Lanceolate |
| ZhongHuang70 | ZH70                     | 2013            | 102             | Zhonghuang13 | Ludou11           | Ovoid      |
| ZhongHuang74 | ZH74                     | 2013            | 109             | Zhongdou27   | Zhonghuang3       | Ovoid      |
| ZhongHuang75 | ZH75                     | 2014            | 131             | NF58         | Teifeng31         | Ovoid      |
| ZhongHuang79 | ZH79                     | 2015            | 131             | Zhongpin 661 | Yudou25           | Ovoid      |

time of the first trifoliate leaf emergence. EDU was repeatedly applied at bi-weekly intervals until the end of the experiment. In total, EDU was applied 7 times during the growing period.

### 1.3. Gas exchange parameters


Photosynthesis at saturating light ( $A_{sat}$ ) and stomatal conductance ( $g_s$ ) were measured using a portable photosynthesis system (LI-6400, LI-COR Inc., Lincoln, NE, USA). For every cultivar, two fully expanded upper leaves per one plant in each plot were randomly selected on the 23rd of August. All measurements were conducted during 08:30–11:00 on clear days under the following conditions: saturating photosynthetic active radiation (PAR) of 1500  $\mu\text{mol}/(\text{m}^2\cdot\text{sec})$ ,  $\text{CO}_2$  at 400 ppm, leaf temperature at 28°C, and relative humidity in air between 50%–70%.

### 1.4. Photosynthetic pigment

After the photosynthesis measurement, the leaf was sampled for photosynthetic pigment. For every cultivar, two leaflets from two fully expanded leaves per plant were randomly punched, and treated with 2 mL 95% ethanol in the dark for 48 hr at 4°C. Assays for chlorophyll (Chl) *a* and *b* and carotenoid (Car) content were carried out by ultraviolet–visible (UV–VIS) spectrophotometry (Alpha-1506, Lab-Spectrum Instruments Co., Ltd., China) according to the specific absorption coefficients provided by Lichtenthaler (1987).

### 1.5. Antioxidant parameters

Leaves for antioxidant analyses were collected immediately after the photosynthesis measurement. Two fully expanded



**Fig. 1 – The 8-hr (9:00–17:00) mean  $\text{O}_3$  concentrations and AOT 40 during the study period from the 20th of June to the 8th of October, 2015. AOT 40: accumulated  $\text{O}_3$  over an hourly concentration threshold of 40 ppb.**

upper leaves were randomly sampled from two plants per each cultivar in each plot, frozen immediately in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  until analysis. Malondialdehyde (MDA), which is related with the level of lipid peroxidation (Feng et al., 2011), was assessed for estimation of lipid peroxidation by 2-thiobarbituric acid-reactive metabolite (TBA) according to the method of Heath and Parker (1968). The optical density (OD) values were obtained in a 96-well plate reader (SpectraMAX 190, Molecular Devices, Sunnyvale, CA, USA) and the MDA content ( $C_{\text{MDA}}$ , mmol/L) was calculated by the equation  $C_{\text{MDA}} = 6.45 \times (A_{532} - A_{600}) - 0.56 \times A_{450}$  ( $A_{532}$ ,  $A_{600}$  and  $A_{450}$  represent extinction value under 450, 532 and 600 nm wavelength, respectively) in order to rule out the disturbances from non-specific ( $A_{600}$ ) and sugar ( $A_{450}$ ) absorbance. Total ascorbate (AsA), which is the main antioxidant metabolite in the mesophyll (Noctor, 2006), was determined using an  $\alpha$ - $\alpha'$ -bipyridyl-based colorimetric assay for approximately 30 mg of ground leaf tissue in a 96-well plate reader SpectraMAX 190 (Gillespie and Ainsworth, 2007).

Samples (~100 mg) for total antioxidant capacity (TAC), which is often used as a synthetic index of the antioxidant pool in the leaves (Gillespie and Ainsworth, 2007), were added to 2 mL cold 70% (v/v) ethanol and homogenized in darkness, according to Benzie and Strain (1996). The mixture was incubated for 20 min in darkness at  $4^{\circ}\text{C}$ , then centrifuged at 3000 r/min for 20 min. One-hundred milliliters of supernatant was taken for the ferric reducing antioxidant power (FRAP) assay to express TAC as  $\text{Fe}^{3+}$  equivalents (mmol  $\text{Fe}^{2+}$ /g fresh mass), and OD values were also measured by a 96-well plate reader SpectraMAX 190 (49i-PS, Thermo Scientific, USA).

For antioxidant enzymes, samples (~0.1 g fresh leaves) were ground in liquid nitrogen and extracted with 2 mL of 50 mmol/L sodium phosphate buffer (pH 7.0, containing 1% vinyl pyrrolidone). The samples were centrifuged at 13,000 r/min,  $4^{\circ}\text{C}$  for 20 min and the supernatant was collected. All extraction experiments were carried out in an ice bath. The superoxide dismutase (SOD) activity was determined by estimating its ability to inhibit the photochemical reduction of nitro blue tetrazolium (NBT), and the absorbance was read at 560 nm. The amount of enzyme that inhibited the 50% NBT reduction was defined as one unit of SOD activity. The ascorbate peroxidase (APX) activity was assayed by monitoring the change of the absorbance at 290 nm after the enzyme extract was mixed a 50 mmol/L potassium phosphate buffer (pH 7.0) containing 30 mmol/L ascorbate and 10 mmol/L  $\text{H}_2\text{O}_2$  (Nakano and Asada, 1981). The enzyme activity was determined by monitoring the change in OD at 290 nm, and one unit of APX activity was defined as the decrease of 0.01  $\Delta\text{OD}$  (changes of optical density) per minute. The catalase (CAT) activity was determined according to the description by Aebi (1984). Peroxidase (POD) activity was assayed adopting the method of Kar and Choudhuri (1987). The unit of CAT and POD activity was defined as the decrease and increase of 0.01  $\Delta\text{OD}$ /min at 240 and 470 nm, respectively. All measurements were conducted using spectrophotometric methods.

## 1.6. Biomass

Plants were harvested on the 8th of October. The above-ground parts of two plants for each cultivar were collected in each plot. Samples were dried in an oven at  $80^{\circ}\text{C}$  until a constant weight was reached.

## 1.7. Definition of $\text{O}_3$ stress effects

The cultivar-specific sensitivity to  $\text{O}_3$  ( $B_{\text{EFF}}$ ) was estimated by comparing the average biomass ( $B$ ) with and without EDU protection, i.e.,  $B_{\text{EFF}} = (B_{\text{EDU}} - B_{\text{Water}}) \times 100/B_{\text{EDU}}$  ( $B_{\text{EDU}}$  means biomass treated with EDU,  $B_{\text{Water}}$  means biomass treated with water). The same approach was used to estimate the effect of  $\text{O}_3$  on each variable  $X$ .

## 1.8. Statistical analyses

The statistical unit was the single plot, with four plots for EDU and three plots for water treatment. Data were checked for normal distribution and homogeneity of variance. Non-normally distributed data i.e., biomass, AsA, POD, and APX, were log transformed prior to analysis. Data in the figures and tables are not transformed, but rather original data means  $\pm$  SD. Data were subject to two-way analysis of variance (ANOVA) including the effects of EDU, cultivar and their interaction using Statistica 10.0 software (StatSoft, Italy). Student's  $t$  test was used to analyze the effect of EDU within each cultivar. Simple linear correlations were applied to test the relationship of all variables  $X$  ( $X_{\text{EFF}}$ ,  $X_{\text{EDU}}$  and  $X_{\text{Water}}$ ) with  $B_{\text{EFF}}$ . Effects were considered significant if  $p < 0.05$ .

# 2. Results

## 2.1. Ozone pollution

The AOT40 index calculated for the whole experimental period was 28.04 ppm-hr (Fig. 1). Daily 8-hr mean  $\text{O}_3$  concentrations varied from 22 to 140 ppb. The daily average along the growing season was 64.9 ppb. There were 141 hr with concentrations higher than 100 ppb, which happened mostly in July (Table 2).

**Table 2 – Distribution of mean hourly ozone concentrations (9–17 hr) in 10 ppb classes during the study period.**

| Concentration (ppb) | Hours            |      |        |           |                   |
|---------------------|------------------|------|--------|-----------|-------------------|
|                     | June (20th–30th) | July | August | September | October (1st–8th) |
| <30                 | 4                | 31   | 14     | 15        | 17                |
| 31–40               | 16               | 16   | 20     | 31        | 13                |
| 41–50               | 10               | 35   | 48     | 44        | 17                |
| 51–60               | 7                | 18   | 57     | 56        | 12                |
| 61–70               | 5                | 23   | 34     | 27        | 1                 |
| 71–80               | 9                | 26   | 17     | 19        | 0                 |
| 81–90               | 5                | 22   | 13     | 15        | 1                 |
| 91–100              | 11               | 17   | 11     | 9         | 1                 |
| 101–110             | 5                | 10   | 10     | 5         | 1                 |
| 111–120             | 2                | 6    | 12     | 7         | 1                 |
| 121–130             | 6                | 11   | 7      | 5         | 0                 |
| 131–140             | 5                | 12   | 1      | 4         | 0                 |
| 141–150             | 1                | 11   | 3      | 2         | 0                 |
| >150                | 2                | 10   | 1      | 1         | 0                 |
| Sum of hours        | 88               | 248  | 248    | 240       | 64                |



## 2.2. Gas exchange parameters

$A_{\text{sat}}$  of cultivars protected by EDU showed a significant variability and was significantly increased by 40% across all cultivars, relative to non-protected plants (Table 3). EDU significantly increased  $A_{\text{sat}}$  in Zhonghuang13 (ZH13), Zhonghuang20 (ZH20), Zhonghuang48 (ZH48) and Zhonghuang49 (ZH49). Interestingly, constitutional  $g_s$  did not significantly differ across the EDU-treated cultivars. Also, the effect of EDU on  $g_s$  was not significant.

The cultivars differed in their foliar content of photosynthetic pigments, as indicated by the higher contents in Zhonghuang74 (ZH74) and Zhonghuang66 (ZH66), and lower content in Zhonghuang39 (ZH39), Zhonghuang44 (ZH44) and Zhonghuang75 (ZH75) (Table 4). EDU significantly increased Chl *a*, Chl *b*, total Chl and Car contents by 26%, 52%, 30% and 23% across all cultivars, respectively. There was significant interaction between EDU and cultivar, as indicated by a significant EDU-induced increase in some cultivars but not in others.

## 2.3. Antioxidant parameters

The foliar content of antioxidant metabolites was significantly affected by the cultivar, the EDU treatment and their interaction (Table 5). Across all cultivars, MDA and AsA were decreased by 9% and 27%, respectively, while TAC was increased by 31% due to the EDU treatment.

Also, the activity of antioxidant enzymes was significantly affected by the cultivar, the EDU treatment and their interaction (Table 6). Across all cultivars, SOD showed a negligible 3% reduction due to the EDU treatment, while APX, POD and CAT increased by 28%, 15% and 8%, respectively.

## 2.4. Biomass and ozone sensitivity of the cultivars

Biomass at harvest showed a significant variation with the cultivar, the EDU treatment and their interaction (Fig. 2a). EDU significantly increased the biomass in 15 out of 19 cultivars (i.e., with the exception of Zhonghuang69 (ZH69), Zhonghuang43 (ZH43), ZH49 and Jidou12 (J12)), thus a significant interaction between EDU and cultivars was found. On average, the biomass of EDU-treated cultivars was 36% higher than that of water-treated cultivars.

The cultivar-specific  $O_3$  sensitivity, expressed as percent variation of biomass when plants were protected by ethylenediurea ( $B_{\text{EFF}}$ ), showed a remarkable variation among cultivars, with the most tolerant cultivars ZH43, ZH49 and ZH69 showing insignificant changes (<10%) and the most sensitive cultivars ZH42, ZH50 and ZH70 showing large variations (>45%) (Fig. 2b).

When linear correlations were applied to test the effects of all variables ( $X_{\text{EDU}}$ ,  $X_{\text{Water}}$  and  $X_{\text{EFF}}$ ) on  $B_{\text{EFF}}$ ,  $MDA_{\text{Water}}$  and  $APX_{\text{EFF}}$  increased significantly with increasing  $O_3$  sensitivity of

**Table 3 – Light-saturated photosynthesis rate ( $A_{\text{sat}}$ ) and stomatal conductance ( $g_s$ ) (mean  $\pm$  SD) of 19 soybean cultivars exposed to ambient  $O_3$  concentrations (water) or protected by ethylenediurea (EDU), levels of significance showing differences between EDU and water for any cultivar, and two-way analysis of variance (ANOVA) (EDU  $\times$  cultivar) results.**

| Cultivar              | $A_{\text{sat}}$ ( $\mu\text{mol}/(\text{m}^2\cdot\text{sec})$ ) |                 |              | $g_s$ ( $\text{mmol}/(\text{m}^2\cdot\text{sec})$ ) |               |              |
|-----------------------|--|-----------------|--------------|---|---------------|--------------|
|                       | EDU  | Water           | Significance | EDU   | Water         | Significance |
| J12                   | 13.0 $\pm$ 2.37  | 10.1 $\pm$ 0.07 | ns           | 403 $\pm$ 93  | 223 $\pm$ 103 | ns           |
| ZH13                  | 15.9 $\pm$ 2.5   | 8.5 $\pm$ 1.35  | **           | 264 $\pm$ 117                                       | 310 $\pm$ 93  | ns           |
| ZH20                  | 14.2 $\pm$ 0.94  | 10.6 $\pm$ 1.1  | **           | 297 $\pm$ 196                                       | 213 $\pm$ 58  | ns           |
| ZH39                  | 16.2 $\pm$ 5.72  | 12.7 $\pm$ 1.52 | ns           | 332 $\pm$ 192                                       | 244 $\pm$ 159 | ns           |
| ZH40                  | 10.4 $\pm$ 1.85  | 10.3 $\pm$ 3.77 | ns           | 243 $\pm$ 61  | 352 $\pm$ 55a | *            |
| ZH41                  | 16.1 $\pm$ 1.69  | 14 $\pm$ 3.1    | ns           | 368 $\pm$ 85  | 230 $\pm$ 72  | ns           |
| ZH42                  | 19.0 $\pm$ 5.62  | 12.7 $\pm$ 2.92 | ns           | 445 $\pm$ 148                                       | 175 $\pm$ 69  | *            |
| ZH43                  | 12.4 $\pm$ 1.86  | 10.3 $\pm$ 2.45 | ns           | 270 $\pm$ 54  | 327 $\pm$ 24  | ns           |
| ZH44                  | 15.6 $\pm$ 1.66  | 11.7 $\pm$ 1.98 | *            | 305 $\pm$ 65  | 335 $\pm$ 102 | ns           |
| ZH48                  | 13.4 $\pm$ 2.04  | 8.3 $\pm$ 0.55  | **           | 220 $\pm$ 171                                       | 259 $\pm$ 135 | ns           |
| ZH49                  | 17.1 $\pm$ 2.37  | 8.7 $\pm$ 1.5   | **           | 428 $\pm$ 107                                       | 154 $\pm$ 57  | *            |
| ZH50                  | 13.6 $\pm$ 3.84  | 7.6 $\pm$ 1.46  | ns           | 300 $\pm$ 157                                       | 117 $\pm$ 38  | ns           |
| ZH62                  | 15.2 $\pm$ 2.23  | 14.7 $\pm$ 0.7  | ns           | 286 $\pm$ 150                                       | 350 $\pm$ 190 | ns           |
| ZH66                  | 15.6 $\pm$ 5.15  | 15.0 $\pm$ 2.53 | ns           | 319 $\pm$ 200                                       | 484 $\pm$ 251 | ns           |
| ZH69                  | 13.6 $\pm$ 1.23  | 12.7 $\pm$ 1.48 | ns           | 335 $\pm$ 136                                       | 416 $\pm$ 136 | ns           |
| ZH70                  | 15.8 $\pm$ 3.4   | 7.5 $\pm$ 1.05  | *            | 263 $\pm$ 143                                       | 270 $\pm$ 90  | ns           |
| ZH74                  | 20.1 $\pm$ 4.94  | 13.7 $\pm$ 0.63 | ns           | 518 $\pm$ 265                                       | 415 $\pm$ 67  | ns           |
| ZH75                  | 18.4 $\pm$ 4.45  | 10.7 $\pm$ 0.73 | *            | 455 $\pm$ 167                                       | 203 $\pm$ 118 | ns           |
| ZH79                  | 18.6 $\pm$ 3.59  | 10.6 $\pm$ 2.49 | *            | 299 $\pm$ 80  | 218 $\pm$ 137 | ns           |
| EDU                   | <0.0001 <sup>a</sup>   |                 |              | 0.590 ns  |               |              |
| Cultivar              | <0.0001 <sup>a</sup>   |                 |              | 0.150 ns  |               |              |
| EDU $\times$ Cultivar | 0.054 ns   |                 |              | 0.043 <sup>*</sup>                                  |               |              |

ns: non-significant; SD: standard deviation.

\*  $p \leq 0.05$ .

\*\*  $p \leq 0.01$ .

<sup>a</sup>  $p \leq 0.001$ .

**Table 4 – Photosynthetic pigments (chlorophyll *a*, *b* (Chl *a*, *b*) and total carotenoids (Car)) (mean  $\pm$  SD) of 19 soybean cultivars exposed to ambient O<sub>3</sub> concentrations (water) or protected by ethylenedurea (EDU), levels of significance showing differences between EDU and water for any cultivar, and two-way ANOVA (EDU  $\times$  cultivar) results.**

| Cultivar              | Chl <i>a</i> (mg/m <sup>2</sup> ) |                 |              | Chl <i>b</i> (mg/m <sup>2</sup> ) |                 |              | Chl <i>a</i> + <i>b</i> (mg/m <sup>2</sup> ) |                 |              | Car (mg/m <sup>2</sup> ) |                 |              |
|-----------------------|-----------------------------------|-----------------|--------------|-----------------------------------|-----------------|--------------|--|-----------------|--------------|--------------------------|-----------------|--------------|
|                       | EDU                               | Water           | Significance | EDU                               | Water           | Significance | EDU  | Water           | Significance | EDU                      | Water           | Significance |
| J12                   | 35.7 $\pm$ 7.41                   | 37.7 $\pm$ 2.6  | ns           | 7.42 $\pm$ 1.69                   | 7.35 $\pm$ 1.23 | ns           | 43.1 $\pm$ 6.46                              | 45 $\pm$ 2.91   | ns           | 4.04 $\pm$ 0.6           | 4.11 $\pm$ 0.39 | ns           |
| ZH13                  | 35.1 $\pm$ 3.23                   | 28.5 $\pm$ 4.83 | ns           | 7.17 $\pm$ 1.22                   | 5.21 $\pm$ 2.42 | ns           | 42.3 $\pm$ 3.86                              | 33.7 $\pm$ 7.25 | ns           | 3.82 $\pm$ 0.49          | 3.2 $\pm$ 0.11  | ns           |
| ZH20                  | 34.5 $\pm$ 4.55                   | 16.3 $\pm$ 1.46 | **           | 9.39 $\pm$ 2.42                   | 5.09 $\pm$ 2.98 | ns           | 43.9 $\pm$ 5.3                               | 21.4 $\pm$ 3.92 | **           | 2.93 $\pm$ 0.51          | 1.84 $\pm$ 0.57 | *            |
| ZH39                  | 22.9 $\pm$ 3.38                   | 23.2 $\pm$ 4.85 | ns           | 5.08 $\pm$ 0.68                   | 3.69 $\pm$ 1.65 | ns           | 28 $\pm$ 3.2                                 | 26.9 $\pm$ 5.18 | ns           | 2.43 $\pm$ 0.55          | 2.51 $\pm$ 0.47 | ns           |
| ZH40                  | 28.7 $\pm$ 2.52                   | 26.1 $\pm$ 2.92 | ns           | 6.04 $\pm$ 3.12                   | 5.39 $\pm$ 4.17 | ns           | 34.7 $\pm$ 4.51                              | 31.5 $\pm$ 6.48 | ns           | 3.1 $\pm$ 0.26           | 2.52 $\pm$ 0.22 | *            |
| ZH41                  | 24.1 $\pm$ 5.98                   | 20 $\pm$ 5.39   | ns           | 5.58 $\pm$ 1.55                   | 3.51 $\pm$ 3.98 | ns           | 29.7 $\pm$ 6.37                              | 23.5 $\pm$ 9.2  | ns           | 2.58 $\pm$ 0.31          | 2.44 $\pm$ 0.45 | ns           |
| ZH42                  | 37.8 $\pm$ 4.12                   | 20.8 $\pm$ 3.59 | **           | 9.51 $\pm$ 1.23                   | 2.35 $\pm$ 0.95 | **           | 47.3 $\pm$ 4.92                              | 23.2 $\pm$ 3.78 | **           | 4.22 $\pm$ 0.46          | 2.6 $\pm$ 0.35  | **           |
| ZH43                  | 29.5 $\pm$ 1.6                    | 22.7 $\pm$ 2.04 | **           | 13.39 $\pm$ 5.97                  | 5.23 $\pm$ 1.61 | ns           | 42.9 $\pm$ 7.43                              | 27.9 $\pm$ 3.57 | **           | 3.19 $\pm$ 0.47          | 2.52 $\pm$ 0.26 | ns           |
| ZH44                  | 21 $\pm$ 6.34                     | 20.2 $\pm$ 4.76 | ns           | 8.2 $\pm$ 5.47                    | 8.13 $\pm$ 3.15 | ns           | 29.2 $\pm$ 1.65                              | 28.3 $\pm$ 1.74 | ns           | 2.26 $\pm$ 0.5           | 2.51 $\pm$ 0.36 | ns           |
| ZH48                  | 34.4 $\pm$ 3.49                   | 18.5 $\pm$ 2.27 | **           | 7.79 $\pm$ 0.44                   | 1.56 $\pm$ 0.56 | **           | 42.2 $\pm$ 3.68                              | 20 $\pm$ 2.82   | **           | 3.5 $\pm$ 0.22           | 2.62 $\pm$ 0.03 | **           |
| ZH49                  | 27 $\pm$ 5.23                     | 23 $\pm$ 5.28   | ns           | 6.07 $\pm$ 2.33                   | 6.17 $\pm$ 3.45 | ns           | 33 $\pm$ 6.6                                 | 29.1 $\pm$ 8.67 | ns           | 3.17 $\pm$ 0.25          | 2.4 $\pm$ 0.11  | **           |
| ZH50                  | 29.4 $\pm$ 2.5                    | 28.4 $\pm$ 1.55 | ns           | 6.45 $\pm$ 1.42                   | 3.18 $\pm$ 1.47 | *            | 35.8 $\pm$ 3.76                              | 31.6 $\pm$ 2.95 | ns           | 3.08 $\pm$ 0.28          | 2.42 $\pm$ 0.07 | *            |
| ZH62                  | 26.3 $\pm$ 5.04                   | 21.9 $\pm$ 1.15 | ns           | 4.35 $\pm$ 1.51                   | 4.18 $\pm$ 0.5  | ns           | 30.6 $\pm$ 3.79                              | 26.1 $\pm$ 1.62 | ns           | 3.04 $\pm$ 0.2           | 2.52 $\pm$ 0.17 | *            |
| ZH66                  | 39.4 $\pm$ 3.73                   | 27.6 $\pm$ 4.52 | *            | 10.86 $\pm$ 2.39                  | 8.08 $\pm$ 1.16 | ns           | 50.2 $\pm$ 6.02                              | 35.7 $\pm$ 4.93 | *            | 3.36 $\pm$ 0.5           | 2.25 $\pm$ 0.44 | *            |
| ZH69                  | 26.5 $\pm$ 6.44                   | 24.9 $\pm$ 4.17 | ns           | 4.03 $\pm$ 0.61                   | 4.93 $\pm$ 1.66 | ns           | 30.5 $\pm$ 7.02                              | 29.9 $\pm$ 5.54 | ns           | 3.34 $\pm$ 0.26          | 2.7 $\pm$ 0.18  | *            |
| ZH70                  | 36.4 $\pm$ 2.45                   | 30.1 $\pm$ 5.88 | *            | 9.78 $\pm$ 1.67                   | 8.1 $\pm$ 1.38  | ns           | 46.2 $\pm$ 3.22                              | 38.2 $\pm$ 4.68 | *            | 3.6 $\pm$ 0.12           | 2.32 $\pm$ 0.34 | **           |
| ZH74                  | 41 $\pm$ 3.54                     | 29.5 $\pm$ 1.84 | **           | 8.93 $\pm$ 1.29                   | 3.97 $\pm$ 2.17 | *            | 49.9 $\pm$ 4.74                              | 33.5 $\pm$ 3.92 | **           | 4.18 $\pm$ 0.43          | 3.27 $\pm$ 0.58 | ns           |
| ZH75                  | 28.3 $\pm$ 5.8                    | 23.4 $\pm$ 5.65 | ns           | 8.42 $\pm$ 4.37                   | 3.91 $\pm$ 3.07 | ns           | 36.8 $\pm$ 6.07                              | 27.3 $\pm$ 2.87 | ns           | 3.01 $\pm$ 0.38          | 3.03 $\pm$ 0.38 | ns           |
| ZH79                  | 30.9 $\pm$ 3.67                   | 24.4 $\pm$ 6.76 | ns           | 11.74 $\pm$ 1.79                  | 8.66 $\pm$ 3.64 | ns           | 40.2 $\pm$ 6.66                              | 33.1 $\pm$ 3.13 | ns           | 3.18 $\pm$ 0.41          | 2.46 $\pm$ 0.32 | ns           |
| EDU                   | <0.0001 <sup>a</sup>              |                 |              | <0.0001 <sup>a</sup>              |                 |              | <0.0001 <sup>a</sup>                         |                 |              | <0.0001 <sup>a</sup>     |                 |              |
| Cultivar              | <0.0001 <sup>a</sup>              |                 |              | <0.0001 <sup>a</sup>              |                 |              | <0.0001 <sup>a</sup>                         |                 |              | <0.0001 <sup>a</sup>     |                 |              |
| EDU $\times$ Cultivar | <0.0001 <sup>a</sup>              |                 |              | <0.0001 <sup>a</sup>              |                 |              | <0.0001 <sup>a</sup>                         |                 |              | <0.0001 <sup>a</sup>     |                 |              |

ns: non-significant.

\*  $p \leq 0.05$ .

\*\*  $p \leq 0.01$ .

<sup>a</sup>  $p \leq 0.001$ .

**Table 5 – Foliar content of malondialdehyde (MDA), total antioxidant capacity (TAC) and total ascorbic acid (AsA) (mean  $\pm$  SD) of 19 soybean cultivars exposed to ambient O<sub>3</sub> concentrations (water) or protected by ethylenediurea (EDU), levels of significance showing differences between EDU and water treatments for any cultivar, and two-way ANOVA (EDU  $\times$  cultivar) results.**

| Cultivar              | MDA (nmol/g)         |                 |              | AsA ( $\mu$ mol/g)   |                 |              | TAC ( $\mu$ mol/g)   |                  |              |
|-----------------------|----------------------|-----------------|--------------|----------------------|-----------------|--------------|----------------------|------------------|--------------|
|                       | EDU                  | Water           | Significance | EDU                  | Water           | Significance | EDU                  | Water            | Significance |
| J12                   | 4.28 $\pm$ 0.17      | 3.74 $\pm$ 0.43 | ns           | 5.35 $\pm$ 1.18      | 4.17 $\pm$ 0.48 | ns           | 20.94 $\pm$ 4.36     | 19.76 $\pm$ 4.78 | ns           |
| ZH13                  | 3.84 $\pm$ 0.4       | 5.39 $\pm$ 0.3  | **           | 3.64 $\pm$ 0.38      | 5.85 $\pm$ 1.13 | **           | 18.63 $\pm$ 3.63     | 17.54 $\pm$ 3.19 | ns           |
| ZH20                  | 4.97 $\pm$ 0.61      | 5.3 $\pm$ 1.79  | ns           | 3.76 $\pm$ 8.11      | 8.01 $\pm$ 0.89 | **           | 21.20 $\pm$ 2.62     | 15.28 $\pm$ 2.11 | **           |
| ZH39                  | 4.92 $\pm$ 1         | 7.32 $\pm$ 1.53 | ns           | 3.54 $\pm$ 0.73      | 4.13 $\pm$ 0.91 | ns           | 19.59 $\pm$ 8.79     | 23.80 $\pm$ 8.42 | ns           |
| ZH40                  | 5.06 $\pm$ 0.42      | 5.68 $\pm$ 1.01 | ns           | 2.84 $\pm$ 0.57      | 5.20 $\pm$ 1.04 | **           | 30.66 $\pm$ 0.94     | 14.99 $\pm$ 1.25 | **           |
| ZH41                  | 4.4 $\pm$ 1.07       | 4.41 $\pm$ 0.59 | ns           | 2.79 $\pm$ 0.76      | 5.18 $\pm$ 0.43 | **           | 12.21 $\pm$ 4.14     | 12.41 $\pm$ 4.74 | ns           |
| ZH42                  | 3.81 $\pm$ 0.5       | 5.83 $\pm$ 0.53 | **           | 3.55 $\pm$ 0.50      | 5.29 $\pm$ 1.03 | *            | 23.74 $\pm$ 4.56     | 12.72 $\pm$ 1.24 | **           |
| ZH43                  | 4.51 $\pm$ 0.86      | 4.98 $\pm$ 0.92 | ns           | 7.21 $\pm$ 0.63      | 5.29 $\pm$ 0.50 | **           | 13.70 $\pm$ 2.48     | 21.95 $\pm$ 2.90 | **           |
| ZH44                  | 4.52 $\pm$ 0.71      | 4.54 $\pm$ 1.56 | ns           | 3.48 $\pm$ 0.81      | 5.76 $\pm$ 1.00 | *            | 7.99 $\pm$ 1.21      | 7.70 $\pm$ 0.85  | ns           |
| ZH48                  | 4.22 $\pm$ 1.39      | 4.65 $\pm$ 0.31 | ns           | 3.41 $\pm$ 0.54      | 3.37 $\pm$ 0.12 | ns           | 20.35 $\pm$ 1.65     | 24.78 $\pm$ 7.55 | ns           |
| ZH49                  | 5.19 $\pm$ 0.13      | 2.94 $\pm$ 0.35 | **           | 4.59 $\pm$ 1.18      | 7.34 $\pm$ 0.42 | **           | 21.69 $\pm$ 2.19     | 13.38 $\pm$ 4.63 | ns           |
| ZH50                  | 4.35 $\pm$ 1.15      | 5.09 $\pm$ 0.53 | ns           | 2.64 $\pm$ 0.61      | 5.97 $\pm$ 0.54 | *            | 10.53 $\pm$ 2.81     | 14.06 $\pm$ 2.22 | *            |
| ZH62                  | 4.27 $\pm$ 0.22      | 4.14 $\pm$ 0.42 | ns           | 4.58 $\pm$ 1.25      | 4.58 $\pm$ 0.23 | ns           | 20.72 $\pm$ 2.35     | 15.96 $\pm$ 2.03 | *            |
| ZH66                  | 3.73 $\pm$ 0.7       | 4.63 $\pm$ 0.6  | ns           | 3.52 $\pm$ 0.63      | 6.12 $\pm$ 1.09 | *            | 19.97 $\pm$ 1.84     | 19.12 $\pm$ 6.60 | ns           |
| ZH69                  | 5.21 $\pm$ 1.65      | 4.92 $\pm$ 1.31 | ns           | 3.54 $\pm$ 0.54      | 5.24 $\pm$ 0.90 | *            | 18.73 $\pm$ 8.39     | 14.42 $\pm$ 4.68 | ns           |
| ZH70                  | 4.71 $\pm$ 0.68      | 6.35 $\pm$ 0.51 | *            | 3.20 $\pm$ 0.82      | 4.06 $\pm$ 0.81 | ns           | 26.68 $\pm$ 1.97     | 16.49 $\pm$ 4.60 | **           |
| ZH74                  | 3.77 $\pm$ 1.34      | 2.74 $\pm$ 0.46 | ns           | 2.60 $\pm$ 0.01      | 4.87 $\pm$ 0.55 | *            | 17.55 $\pm$ 2.22     | 11.07 $\pm$ 0.08 | **           |
| ZH75                  | 3.91 $\pm$ 0.81      | 5.5 $\pm$ 0.94  | ns           | 2.35 $\pm$ 0.37      | 2.77 $\pm$ 0.98 | ns           | 22.54 $\pm$ 3.00     | 13.37 $\pm$ 4.84 | *            |
| ZH79                  | 5.25 $\pm$ 0.99      | 5.84 $\pm$ 0.33 | ns           | 4.11 $\pm$ 0.86      | 4.71 $\pm$ 0.96 | ns           | 20.08 $\pm$ 4.65     | 12.77 $\pm$ 0.33 | **           |
| EDU                   | 0.003**              |                 |              | <0.0001 <sup>a</sup> |                 |              | <0.0001 <sup>a</sup> |                  |              |
| Cultivar              | <0.0001 <sup>a</sup> |                 |              | <0.0001 <sup>a</sup> |                 |              | <0.0001 <sup>a</sup> |                  |              |
| EDU $\times$ Cultivar | 0.002**              |                 |              | <0.0001 <sup>a</sup> |                 |              | <0.0001 <sup>a</sup> |                  |              |

ns: non-significant.  
 \*  $p \leq 0.05$ .  
 \*\*  $p \leq 0.01$ .  
<sup>a</sup>  $p \leq 0.001$ .

the cultivars, while  $B_{\text{Water}}$ ,  $MDA_{\text{EFF}}$  and  $AsA_{\text{EDU}}$  significantly decreased (Table 7).

### 3. Discussion

During the growing season of soybean, the daily average ambient O<sub>3</sub> concentration was 65 ppb and AOT40 was 28 ppm-hr. This level exceeds by far the existing critical level of 3 ppm-hr AOT40 over 3 months recommended in Europe for the protection of agricultural crops (CLRTAP, 2015). Modeling studies confirm that AOT40 levels over China may exceed 15 ppm-hr and are projected to further increase until 2020 (Tang et al., 2013, 2014). While O<sub>3</sub> precursor (mainly NO<sub>x</sub>) emissions have been reduced in Europe and the United States over recent years (Paoletti et al., 2014; Lefohn et al., 2017), emissions are still increasing in China at an annual rate of 5% (Feng et al., 2015). The ozone concentration in Beijing, the capital of China, has been continuously increasing, e.g., the daily maximum 8-hr average O<sub>3</sub> concentration increased by 1.14 ppb/year from 2004 to 2015 (Cheng et al., 2016). Overall, O<sub>3</sub> pollution is a serious concern for many Asian countries, for example India (Singh and Agrawal, 2011; Pandey et al., 2014), Pakistan (Ahmad et al., 2013) and Japan (Hoshika et al., 2011). Beijing and its surroundings, where our experiment was carried out, are a hot-spot of O<sub>3</sub> pollution (Yuan et al., 2015; Zhu et al., 2015).

Even though O<sub>3</sub> pollution is such a pressing issue for food security in China (Yin et al., 2009; Miao et al., 2016) and soybean

is a major staple legume crop for Chinese population (FAO, 2014), previous studies were carried out by simulated O<sub>3</sub> exposure under controlled conditions, i.e., in open-top chambers (Zhang et al., 2014; Zhao et al., 2015). While chambers are well suited for mechanistic studies on O<sub>3</sub> impacts, risk assessment and cultivar screening may be affected by artifacts due to modification of the environmental variables (Paoletti, 2007), as demonstrated in the case of soybean (Howell et al., 1979). EDU has been verified as a useful tool to protect crops from O<sub>3</sub> and assess the effects of O<sub>3</sub> on plants under ambient conditions (Singh et al., 2009; Paoletti et al., 2007; Feng et al., 2011; Hoshika et al., 2013a; Carriero et al., 2015; Yuan et al., 2015; Pandey et al., 2015). Our results confirm that EDU is a valid and easy approach for field assessment of ambient O<sub>3</sub> injury to vegetation.

By using the biomass of EDU-protected plants as a proxy of the biomass in a non-O<sub>3</sub>-polluted environment, we were able to rank the relative O<sub>3</sub>-sensitivity of 19 soybean cultivars widely cultivated in China. The most tolerant cultivars showed insignificant changes (<10% variation of biomass when plants were protected by EDU), while the most sensitive cultivars showed large deviations (>45%). It is well known, in fact, that sensitive plants show significant responses to O<sub>3</sub> when treated with EDU, while tolerant plants show limited responses to O<sub>3</sub> (Szantoi et al., 2007; Singh et al., 2009). Such serious variability in the cultivar-specific O<sub>3</sub> sensitivity has been already shown in other species, e.g., wheat (Biswas et al., 2008; Singh et al., 2009), rice (Akhtar et al., 2010) and tomato

**Table 6 – The activity of antioxidative enzymes (APX, SOD, POD and CAT) (mean  $\pm$  SD) of 19 soybean cultivars exposed to ambient O<sub>3</sub> concentrations (water) or protected by ethylenediurea (EDU), levels of significance showing differences between EDU and water treatments for any cultivar, and two-way ANOVA (EDU  $\times$  cultivar) results.**

| Cultivar              | SOD (U/(g FW))       |                 |              | APX (U/(g FW))       |                |              | POD (U/(g FW))       |                  |              | CAT (U/(g FW))       |                  |              |
|-----------------------|----------------------|-----------------|--------------|----------------------|----------------|--------------|----------------------|------------------|--------------|----------------------|------------------|--------------|
|                       | EDU                  | Water           | Significance | EDU                  | Water          | Significance | EDU                  | Water            | Significance | EDU                  | Water            | Significance |
| J12                   | 305 $\pm$ 54.3       | 372 $\pm$ 60.4  | ns           | 333 $\pm$ 31.9       | 159 $\pm$ 9.7  | ns           | 9.94 $\pm$ 0.54      | 8.52 $\pm$ 0.87  | ns           | 1.70 $\pm$ 0.34      | 3.07 $\pm$ 0.129 | **           |
| ZH13                  | 347 $\pm$ 64.5       | 364 $\pm$ 69.5  | ns           | 154 $\pm$ 10.4       | 286 $\pm$ 59.3 | ns           | 6.17 $\pm$ 0.96      | 6.18 $\pm$ 0.34  | ns           | 1.82 $\pm$ 0.35      | 2.18 $\pm$ 0.19  | ns           |
| ZH20                  | 376 $\pm$ 96.5       | 294 $\pm$ 103.9 | ns           | 233 $\pm$ 6.5        | 131 $\pm$ 7.2  | ns           | 13.7 $\pm$ 0.58      | 9.98 $\pm$ 1.22  | ns           | 2.04 $\pm$ 0.07      | 1.37 $\pm$ 0.22  | **           |
| ZH39                  | 401 $\pm$ 89.7       | 395 $\pm$ 13.1  | ns           | 212 $\pm$ 38         | 162 $\pm$ 59.2 | ns           | 7.10 $\pm$ 1.02      | 6.41 $\pm$ 0.49  | ns           | 2.34 $\pm$ 0.15      | 2.70 $\pm$ 0.16  | *            |
| ZH40                  | 412 $\pm$ 26.3       | 396 $\pm$ 68.1  | ns           | 382 $\pm$ 84         | 242 $\pm$ 18.8 | ns           | 8.70 $\pm$ 0.49      | 8.37 $\pm$ 0.087 | ns           | 4.16 $\pm$ 0.18      | 3.13 $\pm$ 0.25  | **           |
| ZH41                  | 231 $\pm$ 50.3       | 216 $\pm$ 31.2  | ns           | 321 $\pm$ 36.8       | 141 $\pm$ 40.5 | ns           | 9.10 $\pm$ 0.52      | 6.26 $\pm$ 0.12  | **           | 3.38 $\pm$ 0.34      | 1.63 $\pm$ 0.36  | **           |
| ZH42                  | 272 $\pm$ 65.2       | 258 $\pm$ 20.4  | ns           | 538 $\pm$ 117.9      | 253 $\pm$ 19.4 | ns           | 7.37 $\pm$ 0.26      | 9.89 $\pm$ 18.39 | ns           | 2.42 $\pm$ 0.38      | 2.26 $\pm$ 0.43  | ns           |
| ZH43                  | 248 $\pm$ 55.6       | 299 $\pm$ 68.7  | ns           | 110 $\pm$ 5.3        | 280 $\pm$ 16.6 | ns           | 5.83 $\pm$ 0.51      | 8.47 $\pm$ 1.27  | *            | 1.66 $\pm$ 0.22      | 1.89 $\pm$ 0.22  | ns           |
| ZH44                  | 436 $\pm$ 36.4       | 397 $\pm$ 78.6  | ns           | 330 $\pm$ 38.3       | 199 $\pm$ 5.8  | ns           | 8.44 $\pm$ 1.3       | 5.93 $\pm$ 1.02  | ns           | 3.36 $\pm$ 0.26      | 1.52 $\pm$ 0.06  | **           |
| ZH48                  | 425 $\pm$ 34.2       | 355 $\pm$ 51.6  | ns           | 148 $\pm$ 38.4       | 154 $\pm$ 21.6 | ns           | 8.33 $\pm$ 0.24      | 5.85 $\pm$ 0.82  | *            | 1.56 $\pm$ 0.16      | 2.87 $\pm$ 0.12  | **           |
| ZH49                  | 249 $\pm$ 43.9       | 281 $\pm$ 15.8  | ns           | 128 $\pm$ 16.3       | 100 $\pm$ 15.9 | ns           | 13.1 $\pm$ 0.46      | 7.85 $\pm$ 0.54  | ns           | 2.54 $\pm$ 0.14      | 2.13 $\pm$ 0.12  | **           |
| ZH50                  | 354 $\pm$ 56.1       | 302 $\pm$ 56.9  | ns           | 109 $\pm$ 6.1        | 102 $\pm$ 4.3  | ns           | 11.3 $\pm$ 0.61      | 4.46 $\pm$ 0.49  | ns           | 1.57 $\pm$ 0.17      | 2.23 $\pm$ 0.20  | *            |
| ZH62                  | 338 $\pm$ 47.1       | 371 $\pm$ 52.3  | ns           | 376 $\pm$ 6.6        | 131 $\pm$ 21.7 | ns           | 8.29 $\pm$ 0.38      | 7.51 $\pm$ 0.29  | ns           | 1.90 $\pm$ 0.21      | 1.24 $\pm$ 0.24  | *            |
| ZH66                  | 386 $\pm$ 40.9       | 349 $\pm$ 48    | ns           | 533 $\pm$ 52.9       | 163 $\pm$ 2.6  | ns           | 13.7 $\pm$ 0.21      | 10.7 $\pm$ 0.95  | **           | 3.36 $\pm$ 0.19      | 3.34 $\pm$ 0.15  | ns           |
| ZH69                  | 416 $\pm$ 71.3       | 398 $\pm$ 37.8  | ns           | 162 $\pm$ 57         | 400 $\pm$ 73.1 | ns           | 6.80 $\pm$ 0.29      | 7.91 $\pm$ 1.40  | ns           | 1.90 $\pm$ 0.069     | 2.46 $\pm$ 0.13  | **           |
| ZH70                  | 371 $\pm$ 78.8       | 331 $\pm$ 65.7  | ns           | 764 $\pm$ 15         | 235 $\pm$ 19.6 | ns           | 6.78 $\pm$ 0.60      | 7.01 $\pm$ 1.14  | ns           | 2.52 $\pm$ 0.28      | 1.69 $\pm$ 0.30  | *            |
| ZH74                  | 313 $\pm$ 117.8      | 378 $\pm$ 30.7  | ns           | 217 $\pm$ 62.4       | 320 $\pm$ 3.2  | *            | 12.1 $\pm$ 0.44      | 6.93 $\pm$ 0.77  | **           | 3.34 $\pm$ 0.31      | 3.42 $\pm$ 0.21  | ns           |
| ZH75                  | 225 $\pm$ 145.1      | 390 $\pm$ 27.7  | ns           | 155 $\pm$ 50.1       | 203 $\pm$ 4.4  | ns           | 10.3 $\pm$ 0.165     | 6.25 $\pm$ 0.58  | **           | 2.48 $\pm$ 0.25      | 2.98 $\pm$ 0.099 | *            |
| ZH79                  | 250 $\pm$ 38.1       | 381 $\pm$ 46.6  | **           | 210 $\pm$ 31.3       | 228 $\pm$ 23   | ns           | 5.70 $\pm$ 0.016     | 11.2 $\pm$ 1.45  | **           | 1.60 $\pm$ 0.17      | 2.36 $\pm$ 0.19  | **           |
| EDU                   | 0.341 ns             |                 |              | <0.0001 <sup>a</sup> |                |              | <0.0001 <sup>a</sup> |                  |              | <0.0001 <sup>a</sup> |                  |              |
| Cultivar              | <0.0001 <sup>a</sup> |                 |              | <0.0001 <sup>a</sup> |                |              | <0.0001 <sup>a</sup> |                  |              | <0.0001 <sup>a</sup> |                  |              |
| EDU $\times$ Cultivar | 0.049 <sup>*</sup>   |                 |              | <0.0001 <sup>a</sup> |                |              | <0.0001 <sup>a</sup> |                  |              | <0.0001 <sup>a</sup> |                  |              |


ns: non-significant; APX: ascorbate peroxidase; SOD: superoxide dismutase; POD: peroxidase; CAT: Catalase; FW: fresh weight; ANOVA: analysis of variance.

\*  $p \leq 0.05$ .

\*\*  $p \leq 0.01$ .

<sup>a</sup>  $p \leq 0.001$ .





**Fig. 2 – Above-ground biomass (mean  $\pm$  SD) of 19 soybean cultivars exposed to ambient  $O_3$  concentrations (water) or protected by ethylenediurea (EDU), levels of significance showing differences between EDU and water treatments for any cultivar, and two-way ANOVA (EDU  $\times$  cultivar) results. Cultivars are sorted (a) according to decreasing biomass and (b) according to increasing ozone sensitivity, expressed as percent variation when protected by ethylenediurea. The inset shows the result of a two-way analysis of variance (ANOVA) (cultivar  $\times$  EDU treatment). \* $p \leq 0.05$ ; \*\* $p \leq 0.01$ ; \*\*\* $p \leq 0.001$ ; ns: non-significant; SD: standard deviation;  $B_{EFF}$ : the cultivar-specific sensitivity to  $O_3$ ; ZH: Zhonghuang; J12: Jidou12.**

(Calvo et al., 2007), as well as in North American soybean (Burkey and Carter, 2009). This knowledge may be used for breeding novel  $O_3$ -tolerant cultivars as an adaptive strategy to  $O_3$  pollution (Teixeira et al., 2011). As yield is a major plant trait for soybean, further studies including yield are recommended. Current conventional genetic improvement efforts screen for high-yielding cultivars and have indirectly selected genotypes with very high sensitivity to  $O_3$ , as demonstrated in wheat (Biswas et al., 2008). Another short-term adaptive

strategy in hot-spot areas may thus be the cultivation of soybean cultivars that are both high-yielding and  $O_3$ -tolerant (Teixeira et al., 2011), such as our cultivar ZH69, while the most  $O_3$ -sensitive cultivars should be excluded.

The sensitivity of plants to  $O_3$  may be influenced by two main factors:  $O_3$  flux controlled by stomata (Fiscus et al., 2005; Gonzalez-Fernandez et al., 2010) and antioxidant capacity determined by antioxidant pools and enzymes (Biswas et al., 2008; Feng et al., 2010; Inada et al., 2012; Su et al., 2017). We

**Table 7 – Regression coefficient and significance of the linear regressions between the effect of ambient ozone on biomass ( $B_{EFF}$ ) of 19 soybean cultivars, i.e., the cultivar sensitivity to ozone, versus a number of physiological and biochemical variables ( $X$ ) measured in plants exposed to ozone ( $X_{Water}$ ) or protected by using ethylenediurea ( $X_{EDU}$ ).**

| Variable                | $X_{EDU}$ | $X_{Water}$ | $X_{EFF}$ |
|-------------------------|-----------|-------------|-----------|
| Biomass                 | 0.366 ns  | −0.662 **   | –         |
| $A_{sat}$               | 0.255 ns  | −0.250 ns   | 0.252 ns  |
| $g_s$                   | −0.230 ns | −0.141 ns   | 0.210 ns  |
| Chl <i>a</i>            | 0.130 ns  | −0.820 ns   | 0.151 ns  |
| Chl <i>b</i>            | 0.009 ns  | −0.110 ns   | 0.286 ns  |
| Chl <i>a</i> + <i>b</i> | 0.095 ns  | −0.105 ns   | 0.158 ns  |
| Car                     | −0.018 ns | −0.234 ns   | 0.136 ns  |
| MDA                     | −0.332 ns | 0.477 *     | −0.617 ** |
| TAC                     | 0.009 ns  | −0.206 ns   | 0.148 ns  |
| AsA                     | −0.468 *  | −0.137 ns   | −0.118 ns |
| SOD                     | 0.263 ns  | 0.003 ns    | 0.263 ns  |
| APX                     | 0.452 ns  | −0.265 ns   | 0.534 *   |
| CAT                     | 0.001 ns  | −0.140 ns   | 0.096 ns  |
| POD                     | −0.160 ns | −0.152 ns   | 0.005 ns  |

The effect of ozone ( $X_{EFF}$ ) was estimated as  $X_{EFF} = (X_{EDU} - X_{Water}) / X_{EDU} \times 100$ .

\*  $p \leq 0.05$ .

\*\*  $p \leq 0.01$ .

assessed the main physiological and biochemical responses to understand which parameters were the most important as predictors of  $O_3$ -sensitivity in these cultivars and whether the sensitivity mechanisms were similar in the different cultivars. All gas exchange, pigment and biomass responses were consistent in a meta-analysis on EDU effects on 15 crop species exposed to  $O_3$  (Feng et al., 2010), even though the magnitude of changes was higher in our experiment likely due to exposure to higher  $O_3$  concentrations. Also, the responses of antioxidant metabolites and enzymes were consistent with results in the literature for other species treated by EDU (Hassan, 2006; Paoletti et al., 2008; Singh et al., 2009; Pandey et al., 2015). Interestingly, our cultivars differed in both constitutional and  $O_3$ -induced levels of all variables but  $g_s$ . Ozone responses may have been affected by the high variability following the stomatal sluggishness that is a typical response to  $O_3$  (Hoshika et al., 2013b, 2014), although a decline of  $g_s$  is a common response to  $O_3$  (Booker et al., 2009). However, no difference in  $g_s$  among cultivars means that this trait cannot be used for selecting  $O_3$ -tolerant cultivars.

An analysis on how the different variables were related to  $O_3$  sensitivity ( $B_{EFF}$ ) across the cultivars showed significant correlations with: MDA in plants exposed to ambient  $O_3$  ( $MDA_{Water}$ ); percent variation of MDA and APX when plants were protected by EDU ( $MDA_{EFF}$  and  $APX_{EFF}$ ); biomass in plants exposed to ambient  $O_3$  ( $B_{Water}$ ); and constitutional content of AsA in plants protected by EDU ( $AsA_{EDU}$ ). MDA is a marker of  $O_3$ -induced lipid peroxidation in the plant membranes (Feng et al., 2011). This is why  $MDA_{Water}$  was higher in the most sensitive cultivars, which did not possess efficient mechanisms of membrane protection from  $O_3$  injury. As a consequence,  $MDA_{EFF}$ , i.e., the difference in lipid peroxidation between

EDU-protected and non-protected plants, was higher in the most sensitive cultivars. A higher  $APX_{EFF}$  in the most sensitive cultivars was likely due to an excess of  $H_2O_2$ , as APX catalyzes the reduction of  $H_2O_2$  by AsA (Chemikova et al., 2000). Such elevated oxidative stress translated into lower  $B_{Water}$  in the sensitive cultivars, which is a well-known impact of  $O_3$  (Feng and Kobayashi, 2009). The only constitutional factor that explained the cultivar tolerance well was  $AsA_{EDU}$ , as it was lower in the most sensitive cultivars when protected by EDU. While it is well known that the direct reaction of  $O_3$  with cell wall ascorbate is a central mechanism of plant tolerance to this pollutant (Plöchl et al., 2000), this is the first proof linking higher intra-specific  $O_3$  tolerance with higher total ascorbate in soybean. This knowledge will help in breeding for more and more  $O_3$ -tolerant cultivars.

## 4. Conclusions

This was the first experimental study to show that ambient  $O_3$  is able to threaten the growth, physiological and biochemical responses of soybean in China. The results suggested that current  $O_3$  pollution is a serious concern for soybean, which highlights the urgent need for policy-making actions protecting this critical staple legume species for food security in China. Fortunately, cultivars showed a considerable variability in their sensitivity to  $O_3$ , which gives guidance to farmers in choosing the best-suited cultivars for local cultivation. Among the most tolerant cultivars, ZH69 also showed excellent biomass productivity and should be tested for the quantity and quality of yield production.

These important results were obtained by applying the antiozonant EDU as a tool for evaluating the  $O_3$  sensitivity of plants. Although preliminary results are encouraging (Agathokleous et al., 2016a), toxic side effects on the food chain by this synthetic chemical cannot yet be excluded, and thus EDU can be recommended only as a scientific tool and not as an  $O_3$  protectant in common agricultural practice.

The cultivars showed some interesting similarities in their responses to  $O_3$ . We thus conclude that higher lipid peroxidation ( $MDA_{Water}$  and  $MDA_{EFF}$ ) and activity of the ascorbate peroxidase enzyme ( $APX_{EFF}$ ) were responses to  $O_3$  injury, which eventually translated into lower biomass production ( $B_{Water}$ ). Rather than these factors, the constitutional level of total ascorbate ( $AsA_{EDU}$ ) in the leaves was the major parameter explaining soybean cultivar sensitivity to  $O_3$ . This result will guide future breeding efforts towards more  $O_3$ -tolerant soybean cultivars in China, while strategies for implementing control measures of regional  $O_3$  pollution are being implemented.

## Acknowledgments

This work was supported by State Key Laboratory of Soil and Sustainable Agriculture (No. Y20160030), the Hundred Talents Program, Chinese Academy of Sciences, Chinese Academy of Sciences President's International Fellowship Initiative (PIFI) for Senior Scientists (Grant Number 2016VBA057), and CNR-CAS bilateral agreement 2017–2019 (Ozone impacts on plant ecosystems in China and Italy).

## REFERENCES

- Aebi, H., 1984. Catalase *in vitro*. *Methods Enzymol.* 105, 121–126.
- Agathokleous, E., Mouzaki-Paxinou, A.C., Saitanis, C.J., Paoletti, E., Manning, W.J., 2016a. The first toxicological study of the antiozonant and research tool ethylene diurea (EDU) using a *Lemna minor* L. bioassay: hints to its mode of action. *Environ. Pollut.* 213, 996–1006.
- Agathokleous, E., Paoletti, E., Saitanis, C.J., Manning, W.J., Shi, C., Koike, T., 2016b. High doses of ethylene diurea (EDU) are not toxic to willow and act as nitrogen fertilizer. *Sci. Total Environ.* 566–567, 841–850.
- Agathokleous, E., Paoletti, E., Saitanis, C.S., Manning, W.J., Sugai, T., Koike, T., 2016c. Impacts of ethylenediurea (EDU) soil drench and foliar spray in *Salix sachalinensis* protection against O<sub>3</sub>-induced injury. *Sci. Total Environ.* 573, 1053–1062.
- Ahmad, M.N., Bükler, P., Khalid, S., Berge, L.V.D., Shah, H.U., Wahid, A., Emberson, L., Power, S.A., Ashmore, M., 2013. Effects of ozone on crops in north-west Pakistan. *Environ. Pollut.* 174, 244–249.
- Ainsworth, E.A., Yendrek, C.R., Sitch, S., Collins, W.J., Emberson, L.D., 2012. The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annu. Rev. Plant Biol.* 63, 637–661.
- Akhtar, N., Yamaguchi, M., Inada, H., Hoshino, D., Kondo, T., Fukami, M., Funada, R., Izuta, T., 2010. Effects of ozone on growth, yield and leaf gas exchange rates of four Bangladeshi cultivars of rice (*Oryza sativa* L.). *Environ. Pollut.* 158, 2970–2976.
- Ariyaphanphitak, W., Chidthaisong, A., Sarobol, E., Bashkin, V.N., Towprayoon, S., 2005. Effects of elevated ozone concentrations on Thai Jasmine rice cultivars (*Oryza sativa* L.). *Water Air Soil Pollut.* 167, 179–200.
- Avnery, S., Mauzerall, D.L., Liu, J., Horowitz, L.W., 2011. Global crop yield reductions due to surface ozone exposure: 2. Year 2030 potential crop production losses and economic damage under two scenarios of O<sub>3</sub> pollution. *Atmos. Environ.* 45, 2297–2309.
- Benzie, I.F.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a measure of ‘antioxidant power’: the FRAP assay. *Anal. Biochem.* 239, 70–76.
- Betzberger, A.M., Yendrek, C.R., Sun, J., Leisner, C.P., Nelson, R.L., Ort, D.R., Ainsworth, E.A., 2012. Ozone exposure response for U.S. soybean cultivars: linear reductions in photosynthetic potential, biomass and yield. *Plant Physiol.* 160, 1827–1839.
- Biswas, D.K., Xu, H., Li, Y.G., Sun, G.Z., Wang, X.Z., Han, X.G., Jiang, G.M., 2008. Genotypic differences in leaf biochemical, physiological and growth responses to ozone in 20 winter wheat cultivars released over the past 60 years. *Glob. Chang. Biol.* 14, 46–59.
- Booker, F., Muntifering, R., Mcgrath, M., Burkey, K., Decoteau, D., Fiscus, E., et al., 2009. The ozone component of global change: potential effects on agricultural and horticultural plant yield, product quality and interactions with invasive species. *J. Integr. Plant Biol.* 51, 337.
- Burkey, K.O., Carter Jr., T.E., 2009. Foliar resistance to ozone injury in the genetic base of U.S. and Canadian soybean and prediction of resistance in descendent cultivars using coefficient of parentage. *Field Crop Res.* 111, 207–217.
- Calvo, E., Martin, C., Sanz, M.J., 2007. Ozone sensitivity differences in five tomato cultivars: visible injury and effects on biomass and fruits. *Water Air Soil Pollut.* 186, 167–181.
- Carnahan, J.E., Jenner, E.L., Wat, E.K.W., 1978. Prevention of ozone injury to plants by a new protectant chemical. *Phytopathology* 68, 1225–1229.
- Carriero, G., Emiliani, G., Giovannelli, A., Hoshika, Y., Manning, W.J., Traversi, M.L., Paoletti, E., 2015. Effects of long-term ambient ozone exposure on biomass and wood traits in poplar treated with ethylenediurea (EDU). *Environ. Pollut.* 206, 575–581.
- Cheng, N.L., Yunting Li, Y.T., Zhang, D.W., Chen, T., Sun, F., Chen, C., Meng, F., 2016. Characteristics of ground ozone concentration over Beijing from 2004 to 2015: trends, transport, and effects of reductions. *Atmos. Chem. Phys. Discuss.* <http://dx.doi.org/10.5194/acp-2016-508>.
- Chernikova, T., Robinson, J.M., Lee, E.H., Mulchi, C.L., 2000. Ozone tolerance and antioxidant enzyme activity in soybean cultivars. *Photosynth. Res.* 64, 15–26.
- CLRTAP, 2015. Manual on methodologies and criteria for modelling and mapping critical loads and levels and air pollution effects, risks and trends. Chapter 3: Mapping Critical Levels for Vegetation. United Nations Economic Commission for Europe (UNECE) Convention on Longrange Transboundary Air Pollution, Geneva (URL: <http://www.icpmapping.org>).
- Elagöz, V., Manning, W.J., 2005. Factors affecting the effects of EDU on growth and yield of field-grown bush beans (*Phaseolus vulgaris* L.), with varying degrees of sensitivity to ozone. *Environ. Pollut.* 136, 385–395.
- FAO, 2013. FAO statistical yearbook 2013: world food and agriculture. <http://www.fao.org/docrep/018/i3107e/i3107e00.htm>.
- FAO, 2014. Crops processed. <http://www.fao.org/faostat/zh/#data/QD>.
- Feng, Z.Z., Kobayashi, K., 2009. Assessing the impacts of current and future concentrations of surface ozone on crop yield with meta-analysis. *Atmos. Environ.* 43, 1510–1519.
- Feng, Z.Z., Wang, S.G., Szantoi, Z., Chen, S., Wang, X.K., 2010. Protection of plants from ambient ozone by applications of ethylenediurea (EDU): a meta-analytic review. *Environ. Pollut.* 158, 3236–3242.
- Feng, Z.Z., Pang, J., Kobayashi, K., Zhu, J., Ort, D.R., 2011. Differential responses in two varieties of winter wheat to elevated ozone concentration under fully open-air field conditions. *Glob. Chang. Biol.* 17, 580–591.
- Feng, Z.Z., Paoletti, E., Bytnerowicz, A., Harmens, H., 2015. Ozone and plants. *Environ. Pollut.* 202, 215–216.
- Fiscus, E.L., Booker, F.L., Burkey, K.O., 2005. Crop responses to ozone: uptake, modes of action, carbon assimilation and partitioning. *Plant Cell Environ.* 28, 997–1011.
- Gillespie, K.M., Ainsworth, E.A., 2007. Measurement of reduced, oxidized and total ascorbate content in plants. *Nat. Protoc.* 2, 871–874.
- Gonzalez-Fernandez, I., Kaminska, A., Dodmani, M., Goumenaki, E., Quarrie, S., Barnes, J.D., 2010. Establishing ozone flux-response relationships for winter wheat: analysis of uncertainties based on data for UK and Polish genotypes. *Atmos. Environ.* 44, 621–630.
- Hassan, I.A., 2006. Physiology and biochemical response of potato (*Solanum tuberosum* L. cv. Kara) to O<sub>3</sub> and antioxidant chemicals: possible roles of antioxidant enzymes. *Ann. Appl. Biol.* 148, 197–206.
- Heath, R.L., Parker, L., 1968. Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.* 125, 189–198.
- Hoshika, Y., Shimizu, Y., Omasa, K., 2011. A comparison between stomatal ozone uptake and AOT40 of deciduous trees in Japan. *iForest* 4, 128–135.
- Hoshika, Y., Pecori, F., Conese, I., Bardelli, T., Marchi, E., Manning, W.J., Badea, O., Paoletti, E., 2013a. Effects of a three-year exposure to ambient ozone on biomass allocation in poplar using ethylenediurea. *Environ. Pollut.* 180, 299–303.
- Hoshika, Y., Omasa, K., Paoletti, E., 2013b. Both ozone exposure and soil water stress are able to induce stomatal sluggishness. *Environ. Exp. Bot.* 88, 19–23.
- Hoshika, Y., Carriero, G., Feng, Z., Zhang, Y., Paoletti, E., 2014. Determinants of stomatal sluggishness in ozone-exposed deciduous tree species. *Sci. Total Environ.* 481, 453–458.
- Howell, R.K., Koch, E.J., Rose Jr., L.P., 1979. Field assessment of air pollution-induced soybean yield losses. *Agron. J.* 71, 285–288.

- Inada, H., Kondo, T., Akhtar, N., Hoshino, D., Yamaguchi, M., Izuta, T., 2012. Relationship between cultivar difference in the sensitivity of net photosynthesis to ozone and reactive oxygen species scavenging system in Japanese winter wheat (*Triticum aestivum*). *Physiol. Plant.* 146, 217–227.
- Kar, R.K., Choudhuri, M.A., 1987. Possible mechanisms of light-induced chlorophyll degradation in senescing leaves of *Hydrilla verticillata*. *Physiol. Plant.* 70, 729–734.
- Lefohn, A.S., Malley, C.S., Simon, H., Wells, B., Xu, X.B., Zhang, L., Wang, T., 2017. Responses of human health and vegetation exposure metrics to changes in ozone concentration distributions in the European Union, United States, and China. *Atmos. Environ.* 152, 123–145.
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* 148, 350–382.
- Manning, W.J., Paoletti, E., Sandermann, H., Ernst, D., 2011. Ethylenediurea (EDU): a research tool for assessment and verification of the effects of ground level ozone on plants under natural conditions. *Environ. Pollut.* 159, 3283–3293.
- Mateosaparić, I., Redondo, C.A., Villanueva Suárez, M.J., Zapatarevilla, M.A., 2008. Soybean, a promising health source. *Nutr. Hosp.* 23, 305–312.
- Miao, W., Huang, X., Song, Y., 2016. An economic assessment of the health effects and crop yield losses caused by air pollution in mainland China. *J. Environ. Sci.* 56, 102–113.
- Mills, G., Buse, A., Gimeno, B., Bermejo, V., Holland, M., Emberson, L., Pleijel, H., 2007. A synthesis of AOT40-based response functions and critical levels of ozone for agricultural and horticultural crops. *Atmos. Environ.* 41, 2630–2643.
- Morgan, P.B., Ainsworth, E.A., Long, S., 2003. How does elevated ozone impact soybean? A meta-analysis of photosynthesis, growth and yield. *Plant Cell Environ.* 26, 1317–1328.
- Nakano, Y., Asada, K., 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 22, 867–880.
- Noctor, G., 2006. Metabolic signaling in defense and stress: the central roles of soluble redox couples. *Plant Cell Environ.* 29, 409–425.
- Osborne, S.A., Mills, G., Hayes, F., Ainsworth, E.A., Büker, P., Emberson, L., 2016. Has the sensitivity of soybean cultivars to ozone pollution increased with time? An analysis of published dose-response data. *Glob. Chang. Biol.* 9, 3097–3111.
- Pandey, A.K., Majumder, B., Keski-Saari, S., Kontunen-Soppela, S., Pandey, V., Oksanen, E., 2014. Differences in responses of two mustard cultivars to ethylenediurea (EDU) at high ambient ozone concentrations in India. *Agric. Ecosyst. Environ.* 196, 158–166.
- Pandey, A.K., Majumder, B., Keski-Saari, S., Kontunen-Soppela, S., Mishra, A., Sahu, N., Pandey, V., Oksanen, E., 2015. Searching for common responsive parameters for ozone tolerance in 18 rice cultivars in India: results from ethylenediurea studies. *Sci. Total Environ.* 532, 230–238.
- Paoletti, E., 2007. Ozone impacts on forests. *CAB Rev.* 2 (13 pp.).
- Paoletti, E., Manning, W.J., Spaziani, F., Tagliaferro, F., 2007. Gravitational infusion of ethylenediurea (EDU) into trunks protected adult European ash trees (*Fraxinus excelsior* L.) from foliar ozone injury. *Environ. Pollut.* 145, 869–873.
- Paoletti, E., Contran, N., Manning, W.J., Castagna, A., Ranieri, A., Tagliaferro, F., 2008. Protection of ash (*Fraxinus excelsior*) trees from ozone injury by ethylenediurea (EDU): roles of biochemical changes and decreased stomatal conductance in enhancement of growth. *Environ. Pollut.* 155, 464–472.
- Paoletti, E., Contran, N., Manning, W.J., Ferrara, A.M., 2009. Use of the antioxidant ethylenediurea (EDU) in Italy: verification of the effects of ambient ozone on crop plants and trees and investigation of EDU's mode of action. *Environ. Pollut.* 157, 1453–1460.
- Paoletti, E., De Marco, A., Beddows, D.C.S., Harrison, R.M., Manning, W.J., 2014. Ozone levels in European and USA cities are increasing more than at rural sites, while peak values are decreasing. *Environ. Pollut.* 192, 295–299.
- Plöchl, M., Lyons, T., Ollerenshaw, J., Barnes, J., 2000. Simulating ozone detoxification in the leaf apoplast through the direct reaction with ascorbate. *Planta* 210, 454–467.
- Rai, R., Agrawal, M., Kumar, C.K., Agrawal, S.B., Emberson, L., Büker, P., 2015. Application of ethylene diurea (EDU) in assessing the response of a tropical soybean cultivar to ambient O<sub>3</sub>: nitrogen metabolism, antioxidants, reproductive development and yield. *Ecotoxicol. Environ. Saf.* 112, 29–38.
- Reinert, R.A., Eason, G., 2000. Genetic control of O<sub>3</sub> sensitivity in a cross between two cultivars of snap bean. *J. Am. Soc. Hortic. Sci.* 125, 222–227.
- Singh, S., Agrawal, S.B., 2011. Cultivar-specific response of soybean (*Glycine max* L.) to ambient and elevated concentrations of ozone under open top chambers. *Water Air Soil Pollut.* 217, 283–302.
- Singh, S., Agrawal, S.B., Agrawal, M., 2009. Differential protection of ethylenediurea (EDU) against ambient ozone for five cultivars of tropical wheat. *Environ. Pollut.* 157, 2359–2367.
- Singh, E., Tiwari, S., Agrawal, M., 2010. Variability in antioxidant and metabolite levels, growth and yield of two soybean varieties: an assessment of anticipated yield losses under projected elevation of ozone. *Agric. Ecosyst. Environ.* 135, 168–177.
- Singh, A., Agrawal, S.B., Shahi, J.P., Agrawal, M., 2014. Investigating the response of tropical maize (*Zea mays* L.) cultivars against elevated levels of O<sub>3</sub> at two developmental stages. *Ecotoxicology* 23, 1447–1463.
- Su, B., Zhou, M., Xu, H., Zhang, X., Li, Y., Su, H., Xiang, B., 2017. Photosynthesis and biochemical responses to elevated O<sub>3</sub> in *Plantago major* and *Sonchus oleraceus* growing in a lowland habitat of northern China. *J. Environ. Sci.* 53, 113–121.
- Sun, J., Feng, Z., Ort, D.R., 2014. Impacts of rising tropospheric ozone on photosynthesis and metabolite levels on field grown soybean. *Plant Sci.* 226, 147–161.
- Szantoi, Z., Chappelka, A.H., Muntifering, R.B., Somers, G.L., 2007. Use of ethylenediurea (EDU) to ameliorate ozone effects on purple coneflower. *Environ. Pollut.* 150, 200–208.
- Tang, H.Y., Takigawa, M., Liu, G., Zhu, J.G., Kobayashi, K., 2013. A projection of ozone-induced wheat production loss in China and India for the years 2000 and 2020 with exposure-based and flux-based approaches. *Glob. Chang. Biol.* 19, 2739–2752.
- Tang, H.Y., Pang, J., Zhang, G.X., Takigawa, M., Liu, G., Zhu, J.G., Kobayashi, K., 2014. Mapping ozone risks for rice in China for years 2000 and 2020 with flux-based and exposure-based doses. *Atmos. Environ.* 86, 74–83.
- Teixeira, E., Fischer, G., van Velthuis, H., van Dingenen, R., Dentener, F., Mills, G., Walter, C., Ewert, F., 2011. Limited potential of crop management for mitigating surface ozone impacts on global food supply. *Atmos. Environ.* 45, 2569–2576.
- Tiwari, S., Agrawal, M., Manning, W.J., 2005. Assessing the impact of ambient ozone on growth and productivity of two cultivars of wheat in India using three rates of application of ethylenediurea (EDU). *Environ. Pollut.* 138, 153–160.
- Verstraeten, W.W., Neu, J.L., Williams, J.E., Bowman, K.W., Worden, J.R., Boersma, K.F., 2015. Rapid increases in tropospheric ozone production and export from China. *Nat. Geosci.* 8, 690–695.
- Wahid, A., Milne, E., Shamsi, S.R.A., Ashmore, M.R., Marshall, F.M., 2001. Effects of oxidants on soybean growth and yield in the Pakistan Punjab. *Environ. Pollut.* 113, 271–280.
- Yin, P., Fang, X., Yun, Y., 2009. Regional differences of vulnerability of food security in China. *J. Geogr. Sci.* 19, 532–544.
- Yuan, X.Y., Calatayud, V., Jiang, L.J., Manning, W.J., Hayes, F., Tian, Y., Feng, Z.Z., 2015. Assessing the effects of ambient ozone in



- China on snap bean genotypes by using ethylenediurea (EDU). *Environ. Pollut.* 205, 199–208.
- Zhang, W.W., Wang, G.H., Liu, X.B., Feng, Z.Z., 2014. Effects of elevated O<sub>3</sub> exposure on seed yield, N concentration and photosynthesis of nine soybean cultivars (*Glycine max* (L.) Merr.) in Northeast China. *Plant Sci.* 226, 172–181.
- Zhao, T.H., Liu, B., Wang, Y., LIU, Y., Zhao, C.R., Yang, X., et al., 2015. Effects of plant endogenous hormones and antioxidant ability in soybean leaves under UV-B and ozone stress. *Acta Ecol. Sin.* 35, 2695–2702.
- Zhu, Z.L., Sun, X.M., Zhao, F., Meixner, F.X., 2015. Ozone concentrations, flux and potential effect on yield during wheat growth in the Northwest-Shandong Plain of China. *J. Environ. Sci.* 34, 1–9.